

Correction

Corrigendum in “Just Relax: Convex Programming Methods for Identifying Sparse Signals in Noise”

Joel A. Tropp, *Member, IEEE*

Abstract—This correspondence closes a gap in the proof of the main lemma from the paper “Just relax: Convex programming methods for identifying sparse signals in noise” (2006).

Index Terms—Algorithms, approximation methods, basis pursuit, convex program, linear regression, optimization methods, orthogonal matching pursuit, sparse representations.

I. INTRODUCTION

The paper [1] studies the minimizers of the nonsmooth convex function

$$L(\mathbf{b}) = \frac{1}{2} \|\mathbf{s} - \Phi \mathbf{b}\|_2^2 + \gamma \|\mathbf{b}\|_1$$

which plays an important role in sparse approximation and compressive sampling. The key result [1, Lemma 6] is a sufficient condition for the support of the minimizer to be contained within a specified index set. This lemma can then be used to study circumstances in which minimizers of L correctly identify the support of a sparse signal \mathbf{s} contaminated with noise.

It has come to the author’s attention that the “proof” of [1, Lemma 6] relies on a claim that holds only for differentiable convex functions. Although the argument requires some amplification, the lemma is true as originally stated. This correspondence provides a complete proof that corrects the error.

A. Notation

All notation is recycled from [1], but we repeat the essential pieces for the convenience of the reader. As usual, $\|\cdot\|_p$ denotes the ℓ_p vector norm with respect to the standard basis. The angle bracket $\langle \cdot, \cdot \rangle$ represents the Hermitian inner product, which is linear in the first variable and conjugate linear in the second variable.

Let Ω be an index set, and consider the linear space \mathbb{C}^Ω of complex-valued vectors with entries indexed by Ω . The standard basis for \mathbb{C}^Ω is the family $\{\mathbf{e}_\omega : \omega \in \Omega\}$, where the vector \mathbf{e}_ω equals one in the component ω and zero in the remaining components.

We study signals that lie in the space \mathbb{C}^d . Consider a family of vectors $\{\varphi_\omega : \omega \in \Omega\} \subset \mathbb{C}^d$, and form a matrix Φ using these vectors as columns. The matrix maps a vector of coefficients in \mathbb{C}^Ω into a signal by the rule

$$\Phi \mathbf{c} = \sum_{\omega \in \Omega} c_\omega \varphi_\omega.$$

Manuscript received September 05, 2008; revised October 13, 2008. Current version published February 04, 2009. This work was supported in part by ONR Award N00014-08-1-0883.

The author is with the Applied and Computational Mathematics, California Institute of Technology, Pasadena, CA 91125-5000 USA (e-mail: jtropp@acm.caltech.edu).

Communicated by J. Romberg, Associate Editor for Signal Processing.

Digital Object Identifier 10.1109/TIT.2008.2009806

The adjoint maps a signal into a coefficient vector by the rule

$$(\Phi^* \mathbf{s})(\omega) = \langle \mathbf{s}, \varphi_\omega \rangle.$$

Given a subset Λ of Ω , we write Φ_Λ for the submatrix of Φ whose columns are listed in Λ . When Φ_Λ has full column rank, the pseudoinverse is defined by

$$\Phi_\Lambda^\dagger = (\Phi_\Lambda^* \Phi_\Lambda)^{-1} \Phi_\Lambda^*.$$

For a fixed signal \mathbf{s} , we define a coefficient vector $\mathbf{c}_\Lambda = \Phi_\Lambda^\dagger \mathbf{s}$ and a signal approximation $\mathbf{a}_\Lambda = \Phi_\Lambda \mathbf{c}_\Lambda$. This approximation \mathbf{a}_Λ can be seen as the orthogonal projection of \mathbf{s} onto the range of Φ_Λ .

Finally, for a convex function f , we write $\partial f(\mathbf{x})$ for the subdifferential of f at the point \mathbf{x} .

B. The Correlation Condition

To establish [1, Lemma 6], the first step is to study minimizers of the function L that are restricted to have fixed support. We state the result without proof, referring the reader to [1, Lemma 5].

Lemma 1 (Restricted Minimizers): Suppose that Φ_Λ has full column rank, and let \mathbf{b}_* minimize the objective function L over all coefficient vectors supported on Λ . A necessary and sufficient condition on such a minimizer is that

$$\mathbf{e}_\Lambda - \mathbf{b}_* = \gamma (\Phi_\Lambda^* \Phi_\Lambda)^{-1} \mathbf{g}, \quad \text{for } \mathbf{g} \in \partial \|\mathbf{b}_*\|_1 \quad (1)$$

Moreover, the minimizer is unique.

The main result provides a sufficient condition under which the restricted minimizer is also the global minimizer of the objective function.

Lemma 2 (Correlation Condition): Suppose that Φ_Λ has full column rank, and let \mathbf{b}_* minimize the function L over all coefficient vectors supported on Λ . Suppose that

$$\|\Phi^*(\mathbf{s} - \mathbf{a}_\Lambda)\|_\infty < \gamma \left[1 - \max_{\omega \notin \Lambda} \left| \langle \Phi_\Lambda^\dagger \varphi_\omega, \mathbf{g} \rangle \right| \right]$$

where \mathbf{g} verifies (1). It follows that \mathbf{b}_* is the unique global minimizer of L .

Together, these two lemmas provide detailed information about the performance of convex programming methods for sparse approximation, as discussed in [1].

II. PROOF OF LEMMA 2

Let \mathbf{b}_* be the unique minimizer of L over coefficient vectors supported on Λ . We develop a sufficient condition under which

$$L(\mathbf{b}_* + \mathbf{h}) - L(\mathbf{b}_*) > 0$$

whenever the norm of the perturbation \mathbf{h} is small enough. Since the objective function L is convex, it follows that \mathbf{b}_* is the unique global minimizer.

Each perturbation admits a unique decomposition

$$\mathbf{h} = \mathbf{u} + \mathbf{v}$$

where $\text{supp}(\mathbf{u}) \subset \Lambda$ and $\text{supp}(\mathbf{v}) \subset \Lambda^c$. Without loss of generality, we may pose some additional constraints. First, we take $\mathbf{v} \neq \mathbf{0}$, since Lemma 1 already addresses the complementary case. We also instate the bound $\|\mathbf{u}\|_\infty \leq \delta$ for a small, positive number δ , which reflects the requirement that the perturbation is tiny.

To begin the calculation, write the perturbed objective function as

$$L(\mathbf{b}_* + \mathbf{h}) = \frac{1}{2} \|\mathbf{s} - \Phi(\mathbf{b}_* + \mathbf{u}) - \Phi\mathbf{v}\|_2^2 + \gamma \|(\mathbf{b}_* + \mathbf{u}) + \mathbf{v}\|_1.$$

Expand the ℓ_2 norm to obtain

$$\begin{aligned} & \|\mathbf{s} - \Phi(\mathbf{b}_* + \mathbf{u}) - \Phi\mathbf{v}\|_2^2 \\ &= \|\mathbf{s} - \Phi(\mathbf{b}_* + \mathbf{u})\|_2^2 + \|\Phi\mathbf{v}\|_2^2 \\ & \quad - 2\text{Re}\langle \mathbf{s} - \Phi\mathbf{b}_*, \Phi\mathbf{v} \rangle + 2\text{Re}\langle \Phi\mathbf{u}, \Phi\mathbf{v} \rangle. \end{aligned}$$

Since the vectors $\mathbf{b}_* + \mathbf{u}$ and \mathbf{v} have disjoint support

$$\|(\mathbf{b}_* + \mathbf{u}) + \mathbf{v}\|_1 = \|\mathbf{b}_* + \mathbf{u}\|_1 + \|\mathbf{v}\|_1.$$

Combine the last three relations, and identify the quantity $L(\mathbf{b}_* + \mathbf{u})$ to reach

$$\begin{aligned} L(\mathbf{b}_* + \mathbf{h}) - L(\mathbf{b}_*) &= L(\mathbf{b}_* + \mathbf{u}) - L(\mathbf{b}_*) \\ & \quad + \frac{1}{2} \|\Phi\mathbf{v}\|_2^2 - \text{Re}\langle \mathbf{s} - \Phi\mathbf{b}_*, \Phi\mathbf{v} \rangle \\ & \quad + \text{Re}\langle \Phi\mathbf{u}, \Phi\mathbf{v} \rangle + \gamma \|\mathbf{v}\|_1. \end{aligned} \quad (2)$$

This identity holds for each perturbation $\mathbf{h} = \mathbf{u} + \mathbf{v}$.

The next step is to develop a lower bound on the right-hand side of (2). Lemma 1 states that \mathbf{b}_* minimizes L over coefficient vectors supported on Λ . As a result

$$L(\mathbf{b}_* + \mathbf{u}) - L(\mathbf{b}_*) \geq 0.$$

The quadratic term $\|\Phi\mathbf{v}\|_2^2$ is also nonnegative, hence

$$L(\mathbf{b}_* + \mathbf{h}) - L(\mathbf{b}_*) \geq \gamma \|\mathbf{v}\|_1 - |\langle \mathbf{s} - \Phi\mathbf{b}_*, \Phi\mathbf{v} \rangle| - |\langle \Phi\mathbf{u}, \Phi\mathbf{v} \rangle|. \quad (3)$$

It is intuitive that the final term, which is quadratic, has higher order than the other terms, so we will ultimately be able to neglect it.

Let us focus on the second term from the right-hand side of (3). Evidently, we can write

$$\mathbf{v} = \left[\sum_{\omega \notin \Lambda} \theta_\omega \mathbf{e}_\omega \right] \|\mathbf{v}\|_1$$

where $\|\theta\|_1 = 1$. Using this expression, we see that

$$\Phi\mathbf{v} = \left[\sum_{\omega \notin \Lambda} \theta_\omega \varphi_\omega \right] \|\mathbf{v}\|_1.$$

Invoke the triangle inequality and then Jensen's inequality to obtain

$$\begin{aligned} |\langle \mathbf{s} - \Phi\mathbf{b}_*, \Phi\mathbf{v} \rangle| &\leq \left[\sum_{\omega \notin \Lambda} |\theta_\omega| |\langle \mathbf{s} - \Phi\mathbf{b}_*, \varphi_\omega \rangle| \right] \|\mathbf{v}\|_1 \\ &\leq \max_{\omega \notin \Lambda} |\langle \mathbf{s} - \Phi\mathbf{b}_*, \varphi_\omega \rangle| \cdot \|\mathbf{v}\|_1. \end{aligned}$$

To control the third term from the right-hand side of (3), we use standard operator norm bounds. Indeed

$$\begin{aligned} |\langle \Phi\mathbf{u}, \Phi\mathbf{v} \rangle| &= |\langle \Phi^* \Phi\mathbf{u}, \mathbf{v} \rangle| \\ &\leq \|\Phi^* \Phi\mathbf{u}\|_\infty \|\mathbf{v}\|_1 \\ &\leq \delta \|\Phi^* \Phi\|_{\infty, \infty} \|\mathbf{v}\|_1 \end{aligned}$$

where we have applied the bound $\|\mathbf{u}\|_\infty \leq \delta$.

Introduce the last two estimates into (3) to discover that

$$L(\mathbf{b}_* + \mathbf{h}) - L(\mathbf{b}_*) \geq \left[\gamma - \max_{\omega \notin \Lambda} |\langle \mathbf{s} - \Phi\mathbf{b}_*, \varphi_\omega \rangle| - \delta \|\Phi^* \Phi\|_{\infty, \infty} \right] \|\mathbf{v}\|_1.$$

Since we may select δ as small as we like, the right-hand side is strictly positive for each small perturbation \mathbf{h} , provided that

$$\gamma - \max_{\omega \notin \Lambda} |\langle \mathbf{s} - \Phi\mathbf{b}_*, \varphi_\omega \rangle| > 0. \quad (4)$$

The remaining challenge is to find a more desirable condition which ensures that (4) holds.

To that end, we write

$$\mathbf{s} - \Phi\mathbf{b}_* = (\mathbf{s} - \Phi\mathbf{c}_\Lambda) + \Phi(\mathbf{c}_\Lambda - \mathbf{b}_*).$$

By definition, $\Phi\mathbf{c}_\Lambda = \mathbf{a}_\Lambda$. Invoke the fact that $\mathbf{c}_\Lambda - \mathbf{b}_*$ is supported inside Λ along with the characterization from Lemma 1 to see that

$$\Phi(\mathbf{c}_\Lambda - \mathbf{b}_*) = \Phi_\Lambda(\mathbf{c}_\Lambda - \mathbf{b}_*) = \gamma(\Phi_\Lambda^\dagger)^* \mathbf{g}$$

where \mathbf{g} verifies (1). Thus, for each index ω

$$|\langle \mathbf{s} - \Phi\mathbf{b}_*, \varphi_\omega \rangle| \leq |\langle \mathbf{s} - \mathbf{a}_\Lambda, \varphi_\omega \rangle| + \gamma \left| \langle \Phi_\Lambda^\dagger \varphi_\omega, \mathbf{g} \rangle \right|.$$

It follows that a sufficient condition for (4) to hold is that

$$\gamma - \max_{\omega \notin \Lambda} \left[\gamma \left| \langle \Phi_\Lambda^\dagger \varphi_\omega, \mathbf{g} \rangle \right| + |\langle \mathbf{s} - \mathbf{a}_\Lambda, \varphi_\omega \rangle| \right] > 0.$$

This inequality is in force whenever

$$\max_{\omega \notin \Lambda} |\langle \mathbf{s} - \mathbf{a}_\Lambda, \varphi_\omega \rangle| < \gamma \left[1 - \max_{\omega \notin \Lambda} \left| \langle \Phi_\Lambda^\dagger \varphi_\omega, \mathbf{g} \rangle \right| \right]. \quad (5)$$

To complete the argument, we just need to rewrite the left-hand side of the latter relation. By construction, the vector $\mathbf{s} - \mathbf{a}_\Lambda$ is orthogonal to φ_ω for each $\omega \in \Lambda$. Therefore, the left-hand side of (5) does not change if we maximize over all $\omega \in \Omega$

$$\max_{\omega \notin \Lambda} |\langle \mathbf{s} - \mathbf{a}_\Lambda, \varphi_\omega \rangle| = \max_{\omega \in \Omega} |\langle \mathbf{s} - \mathbf{a}_\Lambda, \varphi_\omega \rangle|.$$

Finally, note that

$$\max_{\omega \in \Omega} |\langle \mathbf{s} - \mathbf{a}_\Lambda, \varphi_\omega \rangle| = \|\Phi^*(\mathbf{s} - \mathbf{a}_\Lambda)\|_\infty.$$

We arrive at the sufficient condition stated in Lemma 2.

REFERENCES

- [1] J. A. Tropp, "Just relax: Convex programming methods for identifying sparse signals in noise," *IEEE Trans. Inf. Theory*, vol. 51, no. 3, pp. 1030–1051, Mar. 2006.

Joel A. Tropp (S'03–M'04) received the B.A. degree in Plan II liberal arts honors and the B.S. degree in mathematics, in 1999, graduating *magna cum laude*, and the M.S. and Ph.D. degrees in computational and applied mathematics, in 2001 and in 2004, respectively, all from the University of Texas at Austin. His graduate work was supported by a National Science Foundation (NSF) Graduate Fellowship and a Computational and Applied Mathematics Graduate Fellowship.

In 2004, he joined the Mathematics Department, University of Michigan at Ann Arbor, as Research Assistant Professor. In 2005, he won an NSF Mathematical Sciences Postdoctoral Research Fellowship, and he was appointed T. H. Hildebrandt Research Assistant Professor. Since August 2007, he has been Assistant Professor of Computational and Applied Mathematics at the California Institute of Technology, Pasadena.

Dr. Tropp received an ONR Young Investigator award to support his research on sparse approximation and compressive sampling in March 2008.